

## **Subfossil cladocera from boreal lake gahkozero (The republic of Karelia, Russia) as paleoenvironmental proxies**

Ibragimova A., Frolova L., Dmitriy S.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

### **Abstract**

© SGEM2017. All Rights Reserved. This investigation is aimed to determine general development patterns of the natural and climatic situations and lakes of the boreal zone in the later and postglacial time in the southeastern periphery of the Fennoscandian crystalline shield since the time of the last deglaciation. The analysis of the subfossil Cladoceran community was carried out on the basis of the 23 samples of Lake Gahkozero' bottom sediments in the Republic of Karelia. In the subfossil Cladoceran community of the lake 42 taxa have been identified. The species inhabiting the zones of the Palaearctic and Holarctic are predominant in the lake; most of the identified subfossil remains relate to the pelagic species inhabiting the open part of the reservoir. The discovered subfossil remains of the phytophilous species (representatives of the genera *Alona*, *Alonella*, *Pleuroxus*) indicate the presence of the well-developed zone of macrophytes in the lake. The history of the development and the evolution of the lake is traced by the change of the taxonomic composition in the column of the bottom sediments. In the samples there is a dominance of two taxa - *Bosmina* (*Eubosmina*) cf. *longispina* и *Chydorus* cf. *sphaericus*, which replace each other during the evolution of the lake. The analysis of the variation in the diversity of the biotic groups is carried out using indices, determining the degree of species richness, diversity, and the dominance of Cladoceran communities. Results of statistical and stratigraphic analyses are presented.

<http://dx.doi.org/10.5593/sgem2017/41/S19.075>

---

### **Keywords**

Gahkozero lake, Paleoclimate reconstruction, Republic of Karelia, Subfossil Cladocera

### **References**

- [1] Sleptsov A. M., Klimenko V. V., Synthesis of paleoclimatic data, Nature and society, History and modernity, issue 1, p.118-135, 2005.
- [2] Kalesnik S. V. Soviet Union: geographical description: in 22 vol. Part Uzbekistan. Akramov Z. M., Kalesnik S. V., Eds., Moscow: Thought, 1967, 318 p.
- [3] Subetto D. A. Bottom sediments of lakes: paleolimnological reconstruction. SPb.: The Herzen state pedagogical University of A. I. Herzen, 2009, 309 p.

- [4] Frolova L.A., Subfossil Cladocera (Branchiopoda, Crustacea) in climatic and palaeoenvironmental investigations in Eastern Siberia, International Multidisciplinary Scientific GeoConference, SGEM 2016, Russia, Book 4, Energy and Clean Technologies, Conference Proceedings, vol. 2, 2016, pp. 601-607.
- [5] Frey D.G. The ecological significance of cladoceran remains in lake sediments. *Ecology* 41, 1960, pp. 684-699.
- [6] Frolova L., Ibragimova A., Fedorova I. Stratigraphy of Cladocera in a core from A Yamal Peninsula lake (Arctic Russia), 16th International Multidisciplinary Scientific Geo-Conference SGEM 2016, Conference Proceedings, Book 4, vol. 2, 579-587 pp.
- [7] Ibragimova A.G., Frolova L.A., Grekov I.M. Results of subfossil Cladocera (Branchiopoda, Crustacea) analyses from bottom deposits of Lake Antyukh-Lambina (Kola Peninsula, Murmansk region), *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 2016, vol. 7, №6, pp. 3201-3206.
- [8] Lindholm T., Jakovlev J., Kravchenko A. Zaonezhye Peninsula – The pearl of Lake Onega in Reports of the Finnish environment institute, Biogeography, landscapes, ecosystems and species of Zaonezhye Peninsula, in Lake Onega, Russian Karelia, Tapio Lindholm, Jevgeni Jakovlev & Alexey Kravchenko (eds.), Helsinki: Finnish Environment Institute, Natural Environment Centre, 2014, vol. 40, 364 p.
- [9] Golubev A., Rychanchik D., Romashkin A., Polin A. Geological description of Zaonezhye Peninsula in Reports of the Finnish environment institute, Biogeography, landscapes, ecosystems and species of Zaonezhye Peninsula in Lake Onega, Russian Karelia, Tapio Lindholm, Jevgeni Jakovlev & Alexey Kravchenko (eds.), Helsinki: Finnish Environment Institute, Natural Environment Centre, 2014, vol. 40, 364 p.
- [10] Kulikova V. Biosphere reserve Zaonezhye, Karelian ecological newspaper “Green leaf”, vol. 7, issue 93, p. 3-4, 2008.
- [11] Materials of comprehensive environmental survey underpinning the organization of the State natural Park “Zaonezhye”, 2013, 108 p.
- [12] Demidov I. N., Degradation of the late glaciation in the basin of Onega lake, *Geology and minerals of Karelia*, vol. 8, Petrozavodsk: Karelian Research Centre of Russian Academy of Sciences, 2005, pp. 134-142.
- [13] Szeroczyńska K., Sarmaja-Korjonen K., *Atlas of Subfossil Cladocera from Central and Northern Europe*, Friends of the Lower Vistula Society, 2007, 84 pp.
- [14] Kotov A. A., Sinev A. Ju., Glagolev S. M., Smirnov N. N. Cladocera in Identification Key of zooplankton and zoobenthos of European Russia freshwater, vol.1. Zooplankton. Alekseeva V. R., Calolihina S. Ja., Eds., Moscow: Partnership of scientific publications KMK, 2010, pp. 151-276.
- [15] Kudrin A. I. Species structure and spatial distribution of zooplankton communities in conditions of anthropogenic impact (on the example of the Cheboksary reservoir and its tributaries), Assoc. Professor, Diss. 03.02.08. Nizhniy Novgorod, 2016, 244 p.